

Supplement of Atmos. Meas. Tech., 12, 791–809, 2019
<https://doi.org/10.5194/amt-12-791-2019-supplement>
© Author(s) 2019. This work is distributed under
the Creative Commons Attribution 4.0 License.



Supplement of

Joint retrieval of surface reflectance and aerosol properties with continuous variation of the state variables in the solution space – Part 2: application to geostationary and polar-orbiting satellite observations

Marta Luffarelli and Yves Govaerts

Correspondence to: Marta Luffarelli (marta.luffarelli@rayference.eu)

The copyright of individual parts of the supplement might differ from the CC BY 4.0 License.

S1 BHR evaluation against MODIS data







Figure S1 – Density plots of CISAR BHR retrieval from SEVIRI observations compared with MODIS data




(a) BLUE



(b) RED



(c) NIR



(d) SWIR

Figure S2 – Same as S1 but for CISAR retrieval from PROBA-V observations.

S2 CISAR performances in detecting aerosol events




Figure S3 – High AOT event over Murcia (upper panel) and Granada (bottom panel), Spain, over summer 2015. The yellow triangles represent AERONET observations, the blue dots represent CISAR retrieval from SEVIRI (left panel) and PROBA-V (right panel) observations respectively.




Figure S4 – Same as Fig. S3 but over Sede Boker, Israel, over September 2015.